



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

STATISTICAL REPORTS OF MORBIDITY AND MORTALITY, STATES AND CITIES OF THE UNITED STATES (untabulated).

FLORIDA.—Week ended July 16, 1911. Reports from the State board of health show diphtheria present in 1 locality with 1 case, malaria in 6 localities with 19 cases, smallpox in 12 counties with 63 cases, tuberculosis in 8 localities with 15 cases, typhoid fever in 6 localities with 18 cases.

MASSACHUSETTS.—Week ended April 1, 1911. Population of reporting towns, 2,565,623. Total number of deaths from all causes 816, including diphtheria 11, measles 4, scarlet fever 3, tuberculosis 89, and typhoid fever 4.

Week ended April 8, 1911. Population of reporting towns, 2,524,805. Total number of deaths from all causes 834, including diphtheria 10, measles 4, scarlet fever 4, tuberculosis 90, and typhoid fever 2.

Week ended April 15, 1911. Population of reporting towns, 2,565,623. Total number of deaths from all causes 811, including diphtheria 7, measles 9, scarlet fever 2, tuberculosis 83, and typhoid fever 2.

Week ended April 22, 1911. Population of reporting towns, 2,565,623. Total number of deaths from all causes 862, including diphtheria 13, measles 4, scarlet fever 7, tuberculosis 71, and typhoid fever 3.

Week ended April 29, 1911. Population of reporting towns, 2,565,623. Total number of deaths from all causes 769, including diphtheria 10, measles 8, scarlet fever 2, tuberculosis 75, and typhoid fever 2.

Morbidity.—During the 5 weeks ended April 29, 1911, cases of communicable diseases were reported as follows: Diphtheria 544, measles 2,760, scarlet fever 699, typhoid fever, 115, tuberculosis, pulmonary, 656, tuberculosis other than pulmonary 3, cerebrospinal meningitis 23, meningitis other than cerebrospinal 7, whooping cough 451, varicella 199, erysipelas (not notifiable) 6, mumps (not notifiable) 79, ophthalmia neonatorum 112, anterior poliomyelitis 10, trachoma 4, smallpox 1.

NEW YORK.—Month of May, 1911. Population, 9,113,614. Total number of deaths from all causes 12,383, including diphtheria 208, measles 167, scarlet fever 177, tuberculosis 1,538, typhoid fever 78. Cases reported: Diphtheria 2,139, measles 10,011, scarlet fever 3,953, smallpox 34, tuberculosis 3,407, typhoid fever 358.

UTAH.—Month of June, 1911. Population, 373,351. Total number of deaths from all causes 277, including diphtheria 7, measles 3, scarlet fever 5, tuberculosis 12, typhoid fever 2. Cases reported: Diphtheria 40, measles 588, scarlet fever 116, smallpox 187, tuberculosis 1 (incomplete), typhoid fever 27.